



IGR J17503-2636: a new X-ray transient discovered by INTEGRAL and confirmed by Swift

Chenevez, J.; Jaisawal, G.; Kuulkers, E.; Sanchez-Fernandez, C.; Bazzano, A.; Paizis, A.; Bodaghee, A.; Brandt, S.; Lund, N.; Bozzo, E.

Total number of authors:

11

Publication date:

2019

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Chenevez, J., Jaisawal, G., Kuulkers, E., Sanchez-Fernandez, C., Bazzano, A., Paizis, A., Bodaghee, A., Brandt, S., Lund, N., Bozzo, E., & Ferrigno, C. (2019, Aug 14). IGR J17503-2636: a new X-ray transient discovered by INTEGRAL and confirmed by Swift. The Astronomer's telegram No. ATel #11952

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

GCN
IAUCs

Outside

ATel on [Twitter](#) and [Facebook](#)
ATELstream
ATel Community Site

Other

The Astronomer's Telegram

[Post](#) | [Search](#) | [Policies](#)
[Credential](#) | [Feeds](#) | [Email](#)

27 Aug 2019; 08:25 UT

This space for free for your conference.

[\[Previous](#) | [Next](#) | [ADS](#)]

IGR J17503-2636: a new X-ray transient discovered by INTEGRAL and confirmed by Swift

Aテル #11952; *J. Chenevez, G. Jaisawal (DTU Space, Denmark), E. Kuulkers (ESA/ESTEC, The Netherlands), C. Sanchez-Fernandez (ESA/ESAC, Spain), A. Bazzano (INAF/IAPS, Italy), A. Paizis (INAF/IASF-Mi, Italy), A. Bodaghee (GCSU, USA), S. Brandt, N. Lund (DTU Space, Denmark), E. Bozzo, C. Ferrigno (ISDC, Switzerland), on behalf of a larger collaboration*
on 14 Aug 2018; 19:39 UT

Distributed as an Instant Email Notice Transients
Credential Certification: Jerome CHENEVEZ (jerome@space.dtu.dk)

Subjects: X-ray, Transient

Referred to by Aテル #: [11990](#), [11992](#)



The X-ray monitor JEM-X on-board the INTEGRAL satellite has discovered a new X-ray transient source during the recent Galactic Bulge observation performed on August 11th (revolution 1986) from UTC 07:44 to 12:30 (Aテル #[438](#), Kuulkers et al. 2007, A&A 466, 595, see also: <http://integral.esac.esa.int/BULGE>).

The source is detected at 7 σ for an effective exposure time of 17.9 ks on the combined JEM-X mosaics from which we obtain the following fluxes:

12 \pm 2 mCrab ($\approx 2.0 \times 10^{-10}$ erg/cm²/s) between 3-10 keV,

17 \pm 3 mCrab ($\approx 1.9 \times 10^{-10}$ erg/cm²/s) between 10-25 keV.

JEM-X sees the source at the following position:

RA = 267.58°, Dec = -26.61° with an uncertainty of 1.5 arcmin

leading to the new source name: IGR J17503-2636.

The source is not detected by IBIS/ISGRI above 20 keV, and we estimate a 3- σ upper limit about 10 mCrab (20-40 keV).

Thanks to a 1-ks exposure with Swift/XRT in PC mode between August 13th 19:41-20:13 (UTC) we can refine the source position at:

RA = 267.57527° = 17h 50m 18.06s

Dec = -26.60458° = -26d 36m 16.5s.

The error radius is 2.2 arcsec (90% confidence) obtained with the enhanced method using Swift/UVOT provided by the on-line Swift/XRT data product tool (Evans et al. 2009, MNRAS 397, 1177).

The Swift/XRT 0.3-10 keV count rate was increasing between the two snapshots from 0.3 to 0.5 c/s. The average spectrum is highly absorbed with a derived absorption column between (6.2-13.7) $\times 10^{22}$ cm⁻² (for a galactic column density of 1.3×10^{22} cm⁻² in that direction). The observed 0.3-10 keV flux is estimated between (2.0-8.3) $\times 10^{-11}$ erg/cm²/s.

The next INTEGRAL observations of the Galactic Bulge region are currently scheduled on August 17th and will continue at a rough cadence of one-two observations per week until October 20th. Multi-wavelength follow-up observations are encouraged to unveil the nature of IGR J17503-2636. We thank the Swift Team for scheduling the ToO observation with short notice.

- Related

11992 The possible VVV near-infrared counterpart of IGR J17503-2636
11991 Erratum: Chandra Localization of IGR J17503-2636
11990 Chandra Localization of IGR J17503-2636
11952 IGR J17503-2636: a new X-ray transient discovered by INTEGRAL and confirmed by Swift

[[Telegram Index](#)]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

rrutledge@astronomersteleggram.org

dfox@astronomersteleggram.org

mansi@astronomersteleggram.org